

WO 2004/111656 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
23 December 2004 (23.12.2004)


PCT

(10) International Publication Number
WO 2004/111656 A1

- (51) International Patent Classification⁷: G01N 33/86, (33/49) (74) Agent: ZACCO SWEDEN AB; P.O. Box 23101, S-104 35 Stockholm (SE).
- (21) International Application Number: PCT/SE2004/000940
- (22) International Filing Date: 15 June 2004 (15.06.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0301785-2 15 June 2003 (15.06.2003) SE
- (71) Applicant (*for all designated States except US*): ZAFENA AB [SE/SE]; Sjöbacka Fallhagen 1, S-590 76 Vreta Kloster (SE).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): RÅNBY, Mats [SE/SE]; Sjöbacka Fallhagen 1, S-590 76 Vreta Kloster (SE).
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: COAGULATION TESTS AT AMBIENT TEMPERATURE



(57) Abstract: A method of determining prothrombin time (PT) in a whole blood, anti-coagulated blood, blood plasma or anti-coagulated blood plasma sample at an ambient temperature in the range of 15°C to 45°C is described. The method is performed with liquid reagents and the PT, preferably expressed as International Normalized Ratio (INR), is calculated based on said temperature and the clotting time (CT). A test kit is also described for analysis of PT which comprises temperature recording means, and one or several separate sealed vessels containing reagents, optionally in lyophilized form for reconstitution prior to use, for clotting one or more defined volumes of whole blood, anti-coagulated blood, blood plasma or anti-coagulated blood plasma sample, and optionally time registration means and volume determining means.



Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.